

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/16, 47/44	A1	(11) International Publication Number: WO 95/22319 (43) International Publication Date: 24 August 1995 (24.08.95)
(21) International Application Number: PCT/US95/01943 (22) International Filing Date: 14 February 1995 (14.02.95) (30) Priority Data: 08/197,025 16 February 1994 (16.02.94) US (71) Applicant: ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US). (72) Inventors: BRISKIN, Jacqueline, E.; 1212 Wellington Circle, Buffalo Grove, IL 60089 (US). GUPTA, Pramod, K.; 6986 Bennington Drive, Gurnee, IL 60031 (US). LOYD, Claud; 10413 W. Prairie, Beach Park, IL 60087 (US). KOHLER, Robert, W.; 404 N. Martin, Waukegan, IL 60085 (US). SEMLA, Susan, J.; 2705 Lincolnwood, Evanston, IL 60201 (US). (74) Agents: JANSSEN, Jerry, F. et al.; Abbott Laboratories, CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PROCESS FOR PREPARING FINE PARTICLE PHARMACEUTICAL FORMULATIONS		
(57) Abstract A process for preparing fine particle pharmaceutical formulations having improved throughput and producing greater uniformity of particle size comprises adding to the dry components of the formulation prior to the steps of wetting, extrusion and spheronization, an extrusion aid material selected from pharmaceutically acceptable oils and waxes having a drop point between about 15 °C and 115 °C.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Process for Preparing Fine Particle Pharmaceutical Formulations

Technical Field

This invention relates pharmaceutical formulation processes. More
5 particularly, the present invention concerns a process for preparing fine particle
pharmaceutical formulations by extrusion/spheronization.

Background of the Invention

Conventional processes for preparing fine particle pharmaceutical
10 formulations by extrusion/spheronization involve the steps of blending the dry
ingredients which make up the formulation, wetting the dry powdered blend,
extruding the resulting wetted blend, and forming the extrudate into fine particles by
spheronization.

Generally, the size of particles produced by the above method is limited to
15 particle sizes ranging above about 0.5 mm. Moreover, the amount of water added in
the wetting step must be carefully controlled. Excess water causes the extrudate in the
extrusion step to take on the consistency of "mud" while too little water causes the
wetted material to rupture the screens of the extrusion equipment. The result is that
without very careful process control of the amount of water added to the formulation
20 in the wetting step, batches may be unacceptable with attendant loss of time and/or
money.

There is thus a need for a convenient, cost-effective and efficient method for
making fine particle pharmaceutical formulations which overcome the disadvantages
inherent in prior art methods.

25

Summary of the Invention

This invention provides a process having improved throughput for preparing
fine particle pharmaceutical formulations which exhibit improved uniformity of
particle size and performance characteristics such as drug release. The process is
30 useful for the preparation of formulations comprising fine particles having particle
sizes ranging between about 0.05 mm and about 1 mm which may be used as a
sprinkle formulation for administering a therapeutically active compound to a patient.
The particles may also be used in suspensions, and as a component of tablets and
capsules.

35

pharmaceutically acceptable oils and waxes having a drop point ranging between about 15°C and 115°C; b) thoroughly blending the dry mixture; c) wetting the mixture resulting from step b) to form a granular mixture of the formulation; d) extruding the granular mixture through a mesh; e) spheronizing the extrudate; and f) drying the fine particles resulting from step e) to form a fine particle formulation.

Detailed Description of the Preferred Embodiments

The present invention provides a method for making fine particulate formulations that can be used to administer therapeutically active compounds to a patient. The process has three distinct advantages over prior art processes for making fine particle formulations: first, the amount of wetting agent (e.g. water or water containing one or more additives) added to the blend of dry ingredients in the wetting step does not need to be as carefully controlled; second, the process is capable of producing fine particle formulations in which the particle size may be less than 0.5 mm; and third, the particle size and the performance characteristics of the fine particle formulations produced is more uniform than that resulting from prior art processes.

The formed particles may contain one or more therapeutically active compounds, including pharmaceutically acceptable salts, esters, amides and prodrugs. The therapeutically active compounds may be any therapeutically active compounds for which oral administration is desired. However, the selected therapeutically active compound should be compatible with the selected extrusion aid material and any excipients. Some examples of therapeutically active compounds that may be used in the present invention include, but are not limited to: α -adrenergic agonists such as clonidine and pseudoephedrine; analgesics such as acetaminophen, aspirin, and ibuprofen; antianginals such as verapamil and nifedipine; antibacterials (antibiotics) such as penicillin, erythromycin, tetracycline, amoxicillin, trimethoprim and clarithromycin; antidepressants such as imipramine; antiinflammatory agents such as indomethacin and zileuton; antimigrane agents such as ergotamine; antineoplastics such as methotrexate and etoposide; antivirals such as acyclovir and zidovudine; calcium channel blockers such as diltiazem and verapamil; cardiotonic agents such as digoxin; expectorants such as guaifenesin; bronchodilators such as theophylline; antihypertensives such as methyl dopa; antihistamines such as diphenhydramine, dextromethorphan, phenyltoloxamine, brompheniramine, and chlorpheniramine; diuretics such as furosemide and hydrochlorothiazide; antiepileptics such as tiagabine, phenytoin sodium, divalproex sodium, trimethadione, and paramethadione; central nervous system stimulators such as caffeine and pemoline; decongestants such as

phenylepinephrine and phenylephrine; inorganic salts such as potassium chloride and calcium carbonate; enzymes such as pancreatic enzyme; and vitamins.

The term "pharmaceutically acceptable salts, esters, amides and prodrugs" as used herein refers to those carboxylate salts, amino acid addition salts, esters, amides and prodrugs of the compounds of the present invention which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of patients without undue toxicity, irritation, allergic response, and the like, as well as the zwitterionic forms, where possible, of the compounds of the invention.

The term "salts" refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared *in situ* during the final isolation and purification of the compounds or by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laureate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts and the like. These may include cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium, and the like, as well as non-toxic ammonium, quaternary ammonium and amine cations including, but not limited to, ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine and the like (cf. S. M. Berge, *et al.*, "Pharmaceutical Salts," J. Pharm. Sci., 66: 1-19 (1977)).

Examples of pharmaceutically acceptable non-toxic esters of the compounds of this invention include C₁ to C₄ alkyl esters wherein the alkyl group is a straight or branched chain. Acceptable esters also include C₅ to C₇ cycloalkyl esters as well as arylalkyl esters such as, but not limited to, benzyl, phenylethyl, phenylpropyl and the like. C₁ to C₄ alkyl esters are preferred. Esters of the compounds of the present invention may be prepared according to conventional methods.

Examples of pharmaceutically acceptable, non-toxic amides of the compounds of this invention include amides derived from ammonia, primary C₁ to C₆ alkyl amines and secondary C₁ to C₆ dialkyl amines wherein the alkyl groups are straight or branched chain. In the case of secondary amines the amine may also be in the form of a 5 or 6 membered heterocycle containing one nitrogen atom. Amides derived from ammonia, C₁ to C₃ alkyl primary amides and C₁ to C₂ dialkyl secondary amides

are preferred. Amides of the compounds of the invention may be prepared according to conventional methods.

The term "pro-drug" refers to compounds that are rapidly transformed *in vivo* to yield the parent compound of the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems", Vol. 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987

Typically the therapeutically active agent is present in the formulation in an amount up to about 90 percent by weight of the entire composition.

The extrusion aid is selected so that it does not detrimentally interact with the therapeutically active compound or any excipients and so that it is biocompatible. Preferably, the extrusion aid is a material having a drop point in the range of 15°C to about 115°C. By the term "drop point" as used throughout this specification and claims is meant the temperature at which the material melts or softens to the point where it forms a drop and falls from the thermometer bulb used to take the drop point measurement.

The particular extrusion aid material or mixture of extrusion aid materials that may be used are also selected with regard to the properties of the therapeutically active compound. For example, if the therapeutically active compound degrades at a certain temperature, the extrusion aid material having a melting or softening temperature below the compound's degradation temperature is preferred. Some examples of suitable extrusion aid materials that may be used in the process of the present invention include, but are not limited to: fats, fatty acid esters, hydrogenated vegetable oils, saturated polyglycolized glycerides of hydrogenated vegetable oils, polyethylene glycol esters of hydrogenated vegetable oils such as Lubritab®, Gelucires® and high molecular weight polyethylene glycols. Other compounds having the above referenced properties, which could be used as extrusion aid materials include, but are not limited to, waxes such as carnauba wax, glyceryl esters of aliphatic acids, especially stearic, palmitic and oleic acids, cocoa butter, phospholipids such as lecithins, and sterols such as cholesterol.

The most preferred extrusion aid materials are Lubritab®, a hydrogenated vegetable oil NF obtainable from the Mendell Co. Carmel, NY, Compritol® 888 ATO, glyceryl behenate NF obtainable from Gattefosse, Westwood, NJ, and wax, preferably micronized wax. Preferably, the extrusion aid material is present in the formulations made by the process of this invention in amounts ranging between about

18, 19, 24, 27-29,
30-33

1 percent by weight to about 75 percent by weight based upon the total weight of the formulation.

Fine particle formulations prepared by the process of the present invention may take the form of so-called "sprinkle" formulations which are packaged in a paper, plastic, or foil cachet or in pull-apart capsules. The patient utilizes the formulation by tearing open the cachet or by pulling apart the capsule and then sprinkles the drug formulation over a food for ingestion. In another application, the fine particles produced by the method of the present invention may be sealed inside a capsule. Alternatively, the fine particles produced by the process of this invention may be dispersed within a convention tablet formulation.

The formulations made by the process of this invention may thus contain ingredients in addition to the active therapeutic agent and the extrusion aid material which are chosen to tailor-make the final formulation for its intended purpose. For example, disintegrating agents such as starch, cross-linked polyvinylpyrrolidone (Crospovidone), cross-linked cellulose (Croscarmellose), and sodium starch glycolate may be added to enhance the fast disintegration and dissolution of the fine particles.

Conversely, for rapidly dissolving drugs, conventional binding agents may be added to the formulations made by the process of this invention to retard too-rapid dissolution. Suitable binder agents include polyvinylpyrrolidone (such as Povidone 30 and Povidone 90), carboxymethyl celluloses, and hydroxymethyl celluloses.

In some cases, fillers such as microcrystalline cellulose and lactose may be added to formulations made by the process of this invention. Microcrystalline cellulose extrudes well and undergoes spheronization well to aid in forming fine particles by the process of this invention and is a preferred filler.

The formulation may also include various excipients which are generally chosen from excipients that are conventionally used in solid compositions, such as tablets. Preferred excipients include lactose, mannitol, microcrystalline cellulose (Avicel®) and vitamin E. The excipients, including fillers, may be present in the composition in an amount of about 75 percent to 90 percent by weight based upon the weight of the formulation.

The particles can also be coated, for example, with an enteric coating or a coating which masks any unpleasant taste of the ingredients of the particulate, including masking the taste of therapeutically active compounds, granulating materials or any excipients. A coating may also be used to provide for the controlled release of the therapeutically active compounds from the fine particulate.

The fine particle produced by the disclosed method have the advantages that they have an improved mouth feel; i.e., the particle do not feel gritty or abrasive to the patient. The surface of the particles are also more uniform than particles produced by prior art methods and therefore are more readily coated, if desired. Similarly, the fine particles produced by the present method have improved chemical stability of the therapeutically active compound.

It is also recognized that there are other applications for which the fine particles of the present invention may be useful. For example, a fine particulate may be useful in agriculture to deliver a therapeutically active compound, a fertilizer, or other agents to plants. It is intended that other such applications which employ the fine particles of the present invention fall within the scope of this invention.

General Process Method

In practicing the process of the present invention, the following general procedure is used. First, all of the dry ingredients are thoroughly blended. On a small scale the dry blending may be carried out in stainless steel bowls. For larger quantities, dry mixing of the ingredients may be conveniently carried out in a twin-shell blender of the Patterson-Kelley type. Planetary mixers, for example a Glen mixer or a Hobart mixer, are also conveniently used.

The resulting dry mixture is then wetted by addition of sufficient wetting fluid (e.g. water) to the dry mixture to obtain a granulated solid having the consistency of damp snow or brown sugar. The wetting step is carried out in batches in mixers of the type described above or, conventional equipment which permits the continuous uniform moisturization of the dry blend.

The granulated powder is then fed by auger to conventional extruding equipment where the solid is extruded at high shear through screens of the appropriate mesh size to form threads of the drug formulation. Typical of extrusion equipment for this step is the Model EXDCS-10 Extruder manufactured by Fuji Paudal Co., Ltd. The product at this stage of the process is in the form of long strands of spaghetti-like drug formulation, with the strands having the diameter of the extrusion mesh.

The strands of drug formulation from the previous step are collected and fed to a spheronizing apparatus, typified by the Marumerizer, manufactured by the Fuji Paudal Co. Ltd. or the CF Granulator manufactured by the Vector Corporation. Microcrystalline cellulose or other excipients may be added to the formulation mixture at this point to dust the material to prevent agglomeration. The spheronizer tumbles

the spaghetti-like strands of drug formulation, breaking them up into spheroids of the general diameter of the strands or smaller.

The spheroids of drug formulation produced in the prior step are then dried in a conventional fluid bed dryer such as that manufactured by Niro, Inc. In the final
5 step, the particles are passed through sieves for sizing. The fine particles may then be incorporated into conventional pharmaceutical formulations as cachets, capsules, or by formulation into tablets or caplets.

Example 1

10 Effect of the Presence or Absence of an Extrusion Aid Material on the Extrusion of Pharmaceutical Formulations and the Uniformity of Particle Size

In this example, several formulations were prepared both with and without an extrusion aid (glyceryl behenate). The various formulations compositions and the results upon extrusion appear in Table 1.

15 Examination of the data in Table 1 shows that when an extrusion aid is included in the formulations prepared by the process of the present invention, smooth extrusion of the formulation follows, even at the very small extrusion mesh size of 0.3 mm. For examples in formulation "1a" which included clarithromycin (a therapeutic agent characteristically difficult to extrude), even though a waxy material,
20 Carbopol, was included in the formulation, the extrusion apparatus screens ruptured immediately after extrusion was begun. In contrast, when the same formulation included the preferred extrusion aid, glyceryl behenate, the extrusion screens flexed but did not rupture. Moreover, greater than 70% of the fine particles which were produced were in the desired range of 40-60 mesh.

25 The drug zileuton is characteristically more easily extruded, but in formulation "1c" which lacked an extrusion aid material, the screens of the extrusion apparatus ruptured after a brief period of successful extrusion. In contrast, in formulations "1d"- "1g" which included hydrogentaed vegetable oil as an extrusion aid material, successful extrusion of the drug formulation was achieved.

30

35

Table 1

Formulation	Composition (% by Weight)	Screen Size	Results
1a	Clarithromycin (57.9) Povidone K90 (7.4) Carbopol (34.7)	0.3 mm	Screens ruptured immediately; no particles obtained
1b	Clarithromycin (43.4) Povidone K90 (5.5) Carbopol (26.0) Hydroxypropyl cellulose (5.0) Glyceryl behenate (10.0) Microcrystalline cellulose (10.0)	0.3 mm	Screens flexed but did not rupture; 70.4% yield of 40-60 mesh particles obtained
1c	Zileuton (95) Povidone K90 (5)	0.3 mm	Much flexing of extrusion screens observed; screens finally ruptured
1d	Zileuton* (82) Povidone K90 (5) Microcrystalline cellulose (10.0) Hydrogenated vegetable oil (3)	0.3 mm	Extrusion screens flexed but did not break
1e	Zileuton* (82) Povidone K90 (5) Microcrystalline cellulose (10.0) Hydrogenated vegetable oil (3)	0.3 mm	Extrusion screens flexed but did not break
1f	Zileuton* (72) Povidone K90 (5) Microcrystalline cellulose (20.0) Hydrogenated vegetable oil (3)	0.3 mm	Extrusion screens flexed but did not break
1g	Zileuton* (72) Povidone K90 (5) Microcrystalline cellulose (20.0) Hydrogenated vegetable oil (3)	0.3 mm	Extrusion screens flexed but did not break
1h	Zileuton* (50) Povidone K30 (5) Sodium starch glycolate (5) Microcrystalline cellulose (40)	0.3 mm	Extrusion screens flexed but did not break
1i	Zileuton* (50) Povidone K30 (5) Sodium starch glycolate (5) Microcrystalline cellulose (30) Compitrol® (10)	0.3 mm	No significant flexing of extrusion screens

* Zileuton is N-hydroxy-N-2-((benzo[b]thien-2-yl)ethyl)urea

Example 2**Effect of the Inclusion of an Extrusion Aid Material on the Role of Wetting Fluid (e.g. Water) in Fine Particle Formulations**

In this example, several fine particle pharmaceutical formulations were prepared using zileuton (i.e. N-hydroxy-N-2-((benzo[b]thien-2-yl)ethyl)urea), excipients, disintegrating agents, binders, water, and the preferred extrusion aid material (glyceryl behenate) in accordance with the present invention. The compositions contained water in amounts ranging between 520 ml per kg of formulation (34.2% by weight) to 760 ml per kg of formulation (43.2% by weight). The formulation compositions are presented in Table 2.

Table 2

Formulation		Composition (% by Weight)						
Component	2a	2b	2c	2d	2e	2f	2g	Mean
Zileuton*	50	50	50	50	50	50	50	
Hydroxypropyl cellulose	5	5	5	5	5	5	5	
Sodium starch glycolate	5	5	5	5	5	5	5	
Glyceryl behenate	5	5	5	5	5	5	5	
Avicel™ PH 101	35	35	35	35	35	35	35	
Water (ml/kg)	520	560	600	640	680	720	760	
% by weight	34.2%	35.9%	37.5%	39.0%	40.5%	41.9	43.2	
% Particles in 30-60 mesh	96.4	96.3	98.5	99.0	97.2	97.6	92.9	96.8
% Drug released in 1 hour	53.6	50.1	47.8	46.4	48.4	47.4	46.6	48.8
% Drug released in 2 hours	75.2	72.9	69.9	68.7	70.5	69.3	70.1	70.9
% Drug released in 4.5 hours	95.7	95.0	93.7	93.1	94.5	93.2	93.7	94.1
Release rate (%/√hr)	59.7	60.6	58.6	57.7	59.2	57.5	58.5	58.8

* Zileuton is N-hydroxy-N-2-((benzo[b]thien-2-yl)ethyl)urea

In each of the formulations presented in Table 2, the compositions were extruded successfully through a 0.5 mm mesh screen to produce fine particle formulations having substantially uniform particle size. The generally accepted teachings in the art are that the amount of wetting fluid (e.g. water) contained in the formulation at the time of extrusion is a critical factor in determining success of extrusion and microparticle spheroid size and shape (cf. L. C. Wan, *et al.*, report in the International Journal of Pharmaceutics, 96: 59-65 (1993)). However, the data shown in Table 2 indicate that when an extrusion aid is incorporated into formulations made in accordance with the process of the present invention, the amount of wetting fluid (e.g. water) employed in the wetting step need not be critically controlled to successfully produce fine particle formulations. The data in Table 2 show that for formulations "2a"- "2g", there is substantial uniformity of the particle size of the batches produced by the method of this invention as well as uniformity in drug release profile. For example, all formulations shown in Table 2 had at least 95% particles in the desired size range of 30-60 mesh. Also, all formulations demonstrated consistent drug release from batch to batch. These data indicate that use of the method of the present invention results in fewer batches being rejected with considerable savings in cost and efficiency of processing.

The foregoing examples have been provided to enable one skilled in the art to more fully understand the invention, but are not intended to be read as limiting the scope of the invention as it is defined by the appended claims.

WE CLAIM:

1. A process for the preparation of fine particle pharmaceutical formulations comprising the steps of
 - a) adding to the dry components of the formulation an extrusion aid material selected from pharmaceutically acceptable oils, and waxes having a drop point ranging between about 15°C and 115°C;
 - b) thoroughly blending the dry mixture;
 - c) wetting the mixture resulting from step b) to form a granular mixture of the formulation;
 - d) extruding the granular mixture through a mesh to form an extrudate;
 - e) spheronizing the extrudate; and
 - f) drying the fine particles resulting from step e) to form a fine particle formulation.
2. The process of Claim 1 wherein said extrusion aid is selected from the group consisting of fats, fatty acid esters, saturated polyglycolized glycerides of hydrogenated vegetable oils, polyethylene glycol esters of hydrogenated vegetable oils, high molecular weight polyethylene glycols, waxes, glyceryl esters of aliphatic acids, cocoa butter, phospholipids and sterols, and mixtures thereof.
3. The process of Claim 1 wherein said extrusion aid is added to the components of the pharmaceutical formulation in an amount ranging between about 1 percent by weight to about 75 percent by weight based upon the total weight of the fine particle formulation.
4. The process of Claim 2 wherein said extrusion aid is selected from the group consisting of glyceryl behenate, hydrogenated vegetable oil, and wax.

5. The process of Claim 4 wherein said extrusion aid is glyceryl behenate.
6. The process of Claim 2 wherein said extrusion aid is wax.
7. The process of Claim 1 wherein said fine particle formulation has a particle size below about 1 mm.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 95/01943

A. CLASSIFICATION OF SUBJECT MATTER

A 61 K 9/16, A 61 K 47/44

According to International Patent Classification (IPC) or to both national classification and IPC 6

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A 61 K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT*

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US, A, 4 755 385 (ETIENNE A. et al.) 05 July 1988 (05.07.88), claim 1; column 8, lines 24-40. --	1-4, 6, 7
X	EP, A, 0 350 701 (FARMA RESA S.r.l.) 17 January 1990 (17.01.90), claims 11,13; examples 2,5. --	1-3, 7
X	EP, A, 0 465 338 (RHONE-POULENC NUTRITION ANIMALE) 08 January 1992 (08.01.92), claims 1,4,5; page 2, lines 23-55; page 3, lines 13-17; example 1; table 5.	1-4, 6

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 April 1995

Date of mailing of the international search report

12.05.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

MAZZUCCO e.h.

INTERNATIONAL SEARCH REPORT

-2-

International Application No

PCT/US 95/01943

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB, A, 757 165 (ABBOTT LABORATORIES) 12 September 1956 (12.09.56), page 4, lines 18-41; page 2, lines 115-123; page 3, lines 9-16. --	1, 2
A	EP, A, 0 204 596 (RHONE-POULENC SANTE) 10 December 1986 (10.12.86), claims 1,3,4; column 3, lines 48-53; table 1, example 7. --	1, 4, 5
A	EP, A, 0 438 359 (RHONE-POULENC RORER SA) 24 July 1991 (24.07.91), claims 1,3,4,8,10; page 4, lines 5,6; page 2, line 44 - page 3, line 24. --	1, 4, 5
A	WO, A, 93/17 667 (TAISHO PHARMACEUTICAL CO., LTD) 16 September 1993 (16.09.93), claims 1-4,6; examples 1-13; page 4, lines 14-17. -----	1-4, 6

ANHANG

ANNEX

ANNEXE

zum internationalen Recherchen-
bericht über die internationale
Patentanmeldung Nr.

to the International Search
Report to the International Patent
Application No.

au rapport de recherche inter-
national relatif à la demande de brevet
international n°

PCT/US 95/01943 SAE 104921

In diesem Anhang sind die Mitglieder
der Patentfamilien der in obenge-
nannten internationalen Recherchenbericht
angeführten Patentdokumente angegeben.
Diese Angaben dienen nur zur Unter-
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family
members relating to the patent documents
cited in the above-mentioned inter-
national search report. The Office is
in no way liable for these particulars
which are given merely for the purpose
of information.

La présente annexe indique les
membres de la famille de brevets
relatifs aux documents de brevets cités
dans le rapport de recherche inter-
national visé ci-dessus. Les renseigne-
ments fournis sont donnés à titre indica-
tif et n'engagent pas la responsabilité
de l'Office.

Im Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
US A 4755385	05-07-88	AT E 63819 AU A1 59876/86 AU B2 586262 CA A1 1271138 DD A5 248057 DE A1 3524572 DE C0 3679463 DK A0 3247/86 DK A 3247/86 DK B 165670 DK C 165670 EP A2 208971 EP A3 208971 EP B1 208971 ES AF 2000269 FI A0 862898 FI A 862898 FI B 85649 FI C 85649 GR A 861762 HK A 797/91 HU A2 42945 HU B 195730 IE B 58951 IL A0 79375 IL A1 79375 JP A2 62012717 JP B4 6088905 KR B1 9310586 NO A0 862772 NO A 862772 NO B 175351 NO C 175351 NZ A 216794 PT A 82949 PT B 82949 SG A 721/91 ZA A 8605105	15-06-91 15-01-87 06-07-89 03-07-90 29-07-87 15-01-87 04-07-91 08-07-86 11-01-87 04-01-93 07-06-93 21-01-87 27-01-88 29-05-91 01-02-88 10-07-86 11-01-87 14-02-92 25-05-92 10-11-86 18-10-91 28-09-87 28-07-88 01-12-93 31-10-86 18-01-90 21-01-87 09-11-94 30-10-93 09-07-86 12-01-87 27-06-94 05-10-94 29-11-88 01-08-86 30-01-89 22-11-91 30-03-88
EP A2 350701		IT U0 8821324 IT U 213826	31-05-88 01-03-90
EP A1 465338	08-01-92	AU A1 79395/91 AU B2 643037 BR A 9102684 CA A1 2045883 FR A1 2663818 FR B1 2663818 HU A0 912201 HU A2 62459 HU B 208785 JP A2 4230318 NZ A 238754 US A 5290560 ZA A 9104955	02-01-92 04-11-93 04-02-92 30-12-91 03-01-92 09-07-93 30-12-91 28-05-93 28-01-94 19-08-92 27-09-93 01-03-94 29-04-92
GB A 757165		keine - none - rien	
EP A1 204596	10-12-86	AT E 45283 AU A1 57224/86 AU B2 579012 CA A1 1266841 DE C0 3664867	15-08-89 13-11-86 10-11-88 20-03-90 14-09-89

			DK A	2172/86	10-11-86
			EP B1	204596	09-08-89
			ES A1	554821	16-04-88
			ES A5	554821	16-05-88
			ES A1	9802206	01-07-88
			FI A0	861923	08-05-86
			FI A	861923	10-11-86
			FR A1	2581541	14-11-86
			FR B1	2581541	20-05-88
			JP A2	61260029	18-11-86
			NZ A	216082	06-01-89
			ZA A	8603431	30-12-86
<hr/>					
EP A1	438359	24-07-91	AT E	103489	15-04-94
			AU A1	69481/91	25-07-91
			AU B2	651566	28-07-94
			CA AA	2034413	20-07-91
			CN A	1054189	04-09-91
			CS A2	9100105	13-08-91
			DE C0	69101493	05-05-94
			DE T2	69101493	04-08-94
			EP B1	438359	30-03-94
			ES T3	2062700	16-12-94
			FI A0	910273	18-01-91
			FI A	910273	20-07-91
			FR A1	2657257	26-07-91
			FR B1	2657257	02-09-94
			HU A0	910178	28-08-91
			IL A0	96984	29-03-92
			JP A2	4212359	03-08-92
			NO A0	910218	18-01-91
			NO A	910218	22-07-91
			NZ A	236820	26-11-91
			PL B1	165719	31-01-95
			PT A	96526	15-10-91
			US A	5188838	23-02-93
			US A	5380532	10-01-95
			ZA A	9100376	27-11-91
<hr/>					
WO A1	9317667	16-09-93	AU A1	36484/93	05-10-93
			EP A1	630233	28-12-94
			FI A	944165	09-09-94
			FI A0	944165	09-09-94
			JP A2	6116138	26-04-94
			NO A	943341	09-09-94
			NO A0	943341	09-09-94
<hr/>					